Brandon J. Erickson, MD
Montanna Casey, ATC
Mackenzie Lindeman, ATC
Terry Lin, PA-C
645 Madison Ave New York, NY
200 White Plains Rd 4th Floor Tarrytown, NY
450 Mamaroneck Ave Suite 200 Harrison, NY
Phone: 914-580-9624
Montanna.Casey@rothmanortho.com
Mackenzie.Lindeman@rothmanortho.com
Terry. Lin@rothmanortho.com
www.brandonericksonmd.com



Quadriceps Tendon / Patellar Tendon Repair Physical Therapy Protocol/Prescription

Patient Name:
Surgery: s/p (LEFT / RIGHT) Quadriceps Tendon Patellar Tendon Repair
Date of Surgery:
Frequency: 1 2 3 4 times/week Duration: 1 2 3 4 5 6 Weeks
Phase I – Maximum Protection (Weeks 0 to 6): Weeks 0-4 ☐ Brace locked in full extension for 6 weeks ☐ 50% weight bearing for 3 weeks. 75% weeks 3-4. Wean off crutches at 4 weeks. ☐ Ice and modalities to reduce pain and inflammation ☐ Patella and patella tendon mobility drills ☐ Range of motion – 0° to 30° knee flexion beginning week 3. Increase 10°-15°/week until full range of motion is achieved.
Weeks 4 to 6: ☐ Full weight bearing ☐ Continue patella/patella tendon mobility
Phase II – Progressive Range of Motion and Early Strengthening (Weeks 6 to 12): Weeks 6 to 8: Full weight bearing Open brace 0°-90° when weight bearing once quad control is adequate Continue with swelling control and patella mobility Gradually progress to full range of motion Begin quadriceps setting Begin multi-plane straight leg raising and closed kinetic chain strengthening program focusing on quality VMO function. Initiate open kinetic chain progressing to closed kinetic chain multi-plane hip strengthening Normalize gait pattern Begin stationary bike Initiate pool program Weeks 8 to 10:
 Wean out of brace □ Continue with patella mobility drills □ Normalize gait pattern □ Restore full ROM □ May begin short arc quadriceps contraction 0-30∞

Brandon J. Erickson, MD Montanna Casey, ATC Mackenzie Lindeman, ATC Terry Lin, PA-C 645 Madison Ave New York, NY 200 White Plains Rd 4th Floor Tarrytown, NY 450 Mamaroneck Ave Suite 200 Harrison, NY Phone: 914-580-9624 Montanna.Casey@rothmanortho.com Mackenzie.Lindeman@rothmanortho.com Terry. Lin@rothmanortho.com www.brandonericksonmd.com ☐ Progress open and closed kinetic chain program from bilateral to unilateral ☐ Increase intensity on stationary bike ☐ Begin treadmill walking program Weeks 10 to 12: ☐ Full ROM ☐ Terminal quadriceps stretching ☐ Advance unilateral open and closed kinetic chain strengthening ☐ Initiate proprioception drills ☐ May introduce elliptical trainer Phase III – Progressive Strengthening (Weeks 12 to 16): Weeks 12 to 16: ☐ Advance open and closed kinetic chain strengthening ☐ Increase intensity on bike, treadmill, and elliptical trainer ☐ Increase difficulty and intensity on proprioception drills ☐ Begin gym strengthening: leg press, hamstring curls, ab/adduction; avoid lunges and knee extensions ☐ Begin multi-directional functional cord program Phase IV – Advanced Strengthening and Functional Drills (Weeks 16 to 20): Weeks 16 to 20: \square May begin leg extensions; 30° to 0° ☐ Begin pool running program advancing to land as tolerated Phase V – Plyometric Drills and Return to Sport Phase (Weeks 20 to 24): Weeks 20 to 24: ☐ Advance gym strengthening ☐ Progress running/sprinting program ☐ Begin multi-directional field/court drills ☐ Begin bilateral progressing to unilateral plyometric drills ☐ Follow-up appointment with physician ☐ Sports test for return to competition Comments: Functional Capacity Evaluation ____Work Hardening/Work Conditioning __X_ Teach HEP Modalities _X__Electric Stimulation ___Ultrasound ___ Iontophoresis ___Phonophoresis ___TENS __X__ Heat before _X__Ice after ___Trigger points massage __X Therapist's discretion

Signature Date

Brandon J. Erickson, MD
Montanna Casey, ATC
Mackenzie Lindeman, ATC
Terry Lin, PA-C
645 Madison Ave New York, NY
200 White Plains Rd 4th Floor Tarrytown, NY
450 Mamaroneck Ave Suite 200 Harrison, NY
Phone: 914-580-9624
Montanna.Casey@rothmanortho.com
Mackenzie.Lindeman@rothmanortho.com
Terry. Lin@rothmanortho.com
www.brandonericksonmd.com

