


Brandon J. Erickson, MD
Montanna Casey, ATC
Mackenzie Lindeman, ATC
Terry Lin, PA-C
645 Madison Ave New York, NY
200 White Plains Rd 4th Floor Tarrytown, NY
450 Mamaroneck Ave Suite 200 Harrison, NY
Phone: 914-580-9624
Montanna.Casey@rothmanortho.com
Mackenzie.Lindeman@rothmanortho.com
Terry.Lin@rothmanortho.com
www.brandonericksonmd.com



Ulnar Collateral Ligament Repair/Reconstruction Discharge Instructions

PAIN MANAGEMENT

- **INTERSCALENE NERVE BLOCK** - A combination of local anesthetics are used to numb your shoulder and arm so your brain will not receive any pain signals during and immediately after surgery. The length of effect varies from person to person, but the block usually provides 8-12 hours of pain relief. You will notice a gradual increase in pain as this begins to wear off at which time you can increase the frequency of the oral pain medications as needed.
- **ORAL PAIN MEDICATIONS**
 - Naproxen 500mg
 - Start by taking 1 tablet by mouth twice daily x 2 weeks
 - This medication is an anti-inflammatory medication.
 - Tylenol 500mg
 - ___ This is a medication for pain.
 - ___ Take one tablet by mouth every four hours **as needed for pain levels 1-3**
 - Tramadol 50mg
 - ___ This is a medication for pain.
 - ___ Take one tablet by mouth every four hours **as needed for pain levels 4-7**
 - Oxycodone 5mg
 - ___ This is a medication for pain.
 - ___ Take one tablet by mouth every four hours **as needed for pain levels 8-10**

SIDE EFFECTS OF THESE MEDICATIONS INCLUDE DROWSINESS AND CONSTIPATION. **YOU SHOULD NOT DRIVE OR DRINK ALCOHOL WHILE TAKING THESE PAIN MEDICATIONS.**

- **OTHER MEDICATIONS**
 - Zofran (ondansetron) 4mg
 - This medication is for nausea.
 - Take this medication by mouth every 8 hours as needed for nausea.
 - Pantoprazole/Omeprazole 40mg
 - This medication is a proton pump inhibitor used to reduce the risk of stomach ulcers while taking a NSAID such as Naproxen.
 - This medication should be taken daily by mouth for two weeks.

ACTIVITY INSTRUCTIONS

- **Splint:** Keep your elbow in the splint. You will keep this on for the first 7-10 days until your post-operative office visit.
 - This splint remains on at all times. When you shower, please keep this dry. You may need to cover the splint with a plastic bag so this does not get wet. It is important to keep the bandaged area clean and dry. We will transition you to a hinged elbow brace at your first postoperative visit.
- **ICE MACHINE**
 - Recommend using **1 hour on, 1 hour off for the first 2 days after surgery while awake**
 - Do not place pad directly on skin – make sure there is a barrier such as a t-shirt or towel
 - After the first 2 days you may use the ice machine as needed for comfort
 - If you did not opt for the ice machine, you may use ice packs on the surgical area for 20-30 minutes every 1-2 hours

WOUND CARE & BATHING

- **BANDAGE**
 - You have a large, bulky bandage and splint on your elbow. Please keep this in place until seen at your post-operative appointment in the office for follow up.
 - At your first post-operative appointment, this bandage will be removed and you will be placed into a hinged elbow brace
 - Please do not get the bandage wet. When showering, use a plastic bag to keep the area clean and dry.
- **BATHING**
 - You may shower. You will need to keep bandages in place. You will need to cover elbow with plastic bag in order to keep this dry.
- **SUTURE REMOVAL**
 - Your incision was closed with absorbable sutures, so no stitches will need to be removed
- Keep an eye on your incisions for **signs of infection**:
 - Excessive drainage that is soaking through your dressing, especially if it is pus-like
 - Redness that is spreading out from the edges of your incision
 - Fever or increased warmth around the surgical area

CONTACTING OUR OFFICE

- When to contact our office immediately:
 - Fever > 101.5°F
 - Excessive bleeding from incision(s)
 - Signs of infection of incision(s)
 - Excruciating pain for which the pain medication is not helping

PHYSICAL THERAPY

- You may schedule your first physical therapy visit for **7-10 days after surgery**
- You may begin the home exercises listed below **tomorrow**. You can remove your sling to perform these exercises.
 - **HAND AND WRIST EXERCISES** (10-15 repetitions per set | 2 sets | 3 times daily)
 - With your arm comfortably supported, gently bend wrist back and forth.
 - Curl the fingers into the palm to make a loose fist and then straighten them out.

